

Antihyperon-Hyperon production in antiproton-proton annihilations with PANDA at FAIR

Michael Papenbrock^{1,a}
on behalf of the PANDA collaboration

¹Dept. of Physics and Astronomy, Uppsala University, Box 516, S-751 20 Uppsala, Sweden

Abstract. Hyperon production is an excellent probe of QCD in the confinement domain, and spin observables are a powerful tool in understanding the underlying physics. For the Ω hyperon, seven polarisation parameters can be extracted from the angular distributions of its decay products with the future PANDA experiment at FAIR. Simulation studies reveal great prospects for strange and single charmed hyperon channels with PANDA. Software tools supporting these investigations are currently under development.

Hyperon production in $\bar{p}p \rightarrow \bar{Y}Y$ reactions gives important insights into strangeness and charm production. In particular, the production of single-, multi-strange and single-charmed hyperons from the light quarks in $\bar{p}p$ implies processes where they are replaced by heavier quarks. For any given process the relevant degrees of freedom are given by its energy scale, which for strangeness production is determined by the mass of the strange quark, $m_s \approx 100$ MeV. This is close to the QCD cut-off, $\Lambda_{QCD} \approx 200$ MeV [1], where the coupling constant α_s grows too large for perturbative QCD to be applicable. Therefore, the production of strange hyperons presents an excellent tool to study QCD in the region below the perturbative regime, *i.e.* the confinement domain, about which very little is known so far. On the other hand, the scale of charm production is governed by the mass of the charm quark, $m_c \approx 1300$ MeV, which is more than ten times larger than m_s . In this region, the strong coupling constant becomes $\alpha \approx 0.3$, just barely enough for a perturbative treatment to be valid. Comparing the production of strange and charmed hyperons could give important insights into the differences in the production mechanism at these two energy scales. The production of hyperons in $\bar{p}p \rightarrow \bar{Y}Y$ is often described by theoretical models based on a constituent quark-gluon picture [2]. In case of strange hyperons, models based on the exchange of one or more t -channel kaons exist as well [3]. There have also been efforts to combine both approaches [4]. In order to discriminate between different models, spin observables are very powerful. For hyperons these can be related to the spin of the individual quarks. The Λ hyperon is often modelled as consisting of a spin 0 ud di-quark and an s -quark carrying the hyperon spin. Since the Λ_c^+ hyperon can be described in a similar way, comparing the spin observables of Λ and Λ_c^+ could give deeper insight into the role of spin degrees of freedom in the production of s - and c -quarks.

The density matrix of a particle of arbitrary spin j is given by

$$\rho = \frac{1}{2j+1} I + \sum_{L=1}^{2j} \frac{2j}{2j+1} \sum_{M=-L}^L Q_M^L r_M^L, \quad (1)$$

^ae-mail: michael.papenbrock@physics.uu.se

Table 1. Estimated production rates of various hyperon channels [8, 15]. The cross sections marked with a * have a very large uncertainty. Details are given in the text.

Beam mom. (GeV/c)	Reaction	σ (μb)	Eff (%)	Decay	Rate at $1 \cdot 10^{31} \text{cm}^{-2} \text{s}^{-1}$
1.64	$\bar{p}p \rightarrow \bar{\Lambda}\Lambda$	64	11	$\Lambda \rightarrow p\pi^-$	29 s^{-1}
4	$\bar{p}p \rightarrow \bar{\Lambda}\Sigma^0$	≈ 40	≈ 30	$\Sigma^0 \rightarrow \Lambda\gamma$	30 s^{-1}
4	$\bar{p}p \rightarrow \bar{\Xi}^+\Xi^-$	≈ 2	≈ 20	$\Xi^- \rightarrow \Lambda\pi^-$	1.5 s^{-1}
12	$\bar{p}p \rightarrow \bar{\Omega}^+\Omega^-$	$\approx 0.002^*$	≈ 30	$\Omega \rightarrow \Lambda K^-$	$\approx 4 \text{ h}^{-1}$
12	$\bar{p}p \rightarrow \bar{\Lambda}_c\Lambda_c^+$	$\approx 0.1^*$	≈ 35	$\Lambda_c \rightarrow \Lambda\pi^+$	$\approx 2 \text{ d}^{-1}$




Figure 2. (a) The polarisation of the Ξ^- as a function of $\cos \theta_{\Xi}$. (b) The spin correlation C_{xx} in the x direction (\hat{m} in Fig. 1) of the $\bar{\Xi}^+$ and the Ξ^- . (c) The polarisation parameter r_0^2 of the Ω as a function of $\cos \theta_{\Omega}$. The lines represent the input sine functions (see text) and the points and bars the reconstructed values for 10000 simulated events.

from 1.5 to 15 GeV/c, interacting with an internal hydrogen target. The PANDA experiment will have nearly full angular coverage with precise tracking and vertex reconstruction, particle identification, calorimetry, and will benefit from the high luminosity of HESR [11]. Simulation studies with a simplified Monte Carlo framework show excellent prospects with high signal and low background rates and a good detection efficiency across the full phase space for all strange and single-charmed hyperon channels [8, 15]. The estimated signal rates in Table 1 are calculated using the day⁻¹ luminosity mode of HESR of $1 \cdot 10^{31} \text{cm}^{-2} \text{s}^{-1}$. The only well known cross section is the one for the $\bar{\Lambda}\Lambda$ channel at 1.64 GeV/c, whereas the cross sections for the other strange channels are extrapolated from data at other energies. For the heavier hyperons, *i.e.* Ω and Λ_c , where no data exist from $\bar{p}p$ annihilation, the values are taken from theoretical predictions which can differ by an order of magnitude [13, 14].

Simulation studies of spin observables for Λ , Σ , Ξ , Ω , and Λ_c hyperons have been performed as well. Since there exist no data on multi-strange or charmed hyperon channels, a parametrisation based on sine functions was introduced at the event generation stage. These fulfil the requirement of the definition of the production plane (see figure 1) of being equal to zero at $\cos \theta_y = \pm 1$. Figure 2 shows some examples of spin observables of Ξ and Ω , parametrised by $P_{y,\Xi} = \sin 2\theta_{\Xi}$, $C_{xx,\Xi} = \sin \theta_{\Xi}$ and $r_{0,\Omega}^2 = \sin 2\theta_{\Omega} / \sqrt{3}$. It is clear that these can be accurately reconstructed with PANDA.

The weak decay of the hyperons goes along with comparatively long lifetimes. Their decay vertices can easily be displaced by several tens centimetres from the production point, setting high demands on precise track reconstruction. The PANDA experiment will include a variety of subdetectors designed for charged track reconstruction, including a Micro Vertex Detector (MVD), a Straw Tube Tracker (STT), and a Gas Electron Multiplier (GEM). The reconstruction can also be supported by

a Scintillator Tile Hodoscope (SciTil). Multiple algorithms for pattern recognition and tracking are currently under development with different algorithms for different detectors or detector groups. A special focus is given here on the STT, which is a large barrel of 4636 straw tubes. Since its main purpose is to gather transverse information from the tracks, most of these are aligned parallel to the beam axis, while a few layers of straws are skewed by $\pm 2.9^\circ$, allowing to gain longitudinal information as well. Simulation studies show that tracks may completely miss one or more of the aforementioned subdetectors, creating only hits within the STT. In these cases, the skewed straws of the STT are the only viable source for a longitudinal momentum information of the track. A tool to support a standalone track finder for the STT is under development which matches the isochrones from the skewed straws with axial information gained from the parallel straws to gather p_z information [16].

Using the high luminosity mode of HESR, the mean time between events can be as small as 100 ns while the drift times in the STT can be up to 250 ns, usually causing hits from multiple events to overlap. Events are reconstructed on a software level, using time gaps between hits in one or more of the faster subdetectors, *e.g.* the MVD. This procedure also yields a first approximation of the start time of each event, T_0 , which in turn is required to determine the isochrones from the straw tube signals. With this first approximation, preliminary tracks will be propagated to detectors with excellent time resolution, *e.g.* the SciTil, in order to acquire a precise T_0 .

References

- [1] Ellis R K, Stirling W J and Webber B R, *QCD and Collider Physics* (Cambridge Monographs on particle physics, Nuclear Physics and Cosmology, Cambridge, 1996)
- [2] Kohno M and Weise W, Phys. Lett. B **179**, 15 (1986); Rubinstein H R and Snellman H, Phys. Lett. B **165**, 187 (1985); Furuji S and Faessler A, Nucl. Phys. A **486**, 669 (1987); Burkardt M and Dillig M, Phys. Rev. C **37**, 1362 (1988); Alberg M A et al., Z. Phys. A **331**, 207 (1988)
- [3] Tabakin F and Eisenstein R A, Phys. Rev. C **31**, 1857 (1985); Kohno M and Weise W, Phys. Lett. B **179**, 15 (1986); La France P et al., Phys. Lett. B **214**, 317 (1988); Timmermanns R G E et al., Phys. Rev. D **45**, 2288 (1992); Haidenbauer J et al., Phys. Rev. C **46**, 2516 (1992)
- [4] Ortega P G et al., Phys. Lett. B **696**, 352 (2011)
- [5] Doncel M G et al., Nucl. Phys B **38**, 477 (1972); Doncel M G et al., Phys. Rev. D **7**, 815 (1973)
- [6] Yao W M et al., J. Phys. G **33**, 1 (2006)
- [7] Koch W ed. Nikolic M, *Analysis of scattering and decay* (Gordon and Breach, New York-London-Paris, 1968)
- [8] Thomé E, *Multi-Strange and Charmed Antihyperon-Hyperon Physics for PANDA* (Ph. D. Thesis, Uppsala University, 2012)
- [9] Johansson T, AIP Conf. Proc. 8th Int. Conf. on Low Energy Antiproton Physics, 95 (2003)
- [10] Klempt E et al., Phys. Rep. **368**, 119 (2002)
- [11] The PANDA collaboration, Technical Performance Report, (2005)
- [12] The PANDA collaboration, Physics Performance Report, 43 (2009)
- [13] Kaidalov A B and Volkovitsky P E, Z. Phys. C **63**, 51 (1994)
- [14] Titov A I and Kämpfer B 2008, Phys. Rev. C **78**, 025201 (2008); Gornitschnig A T et al., Eur. Phys. J A **42**, 43 (2009); He J et al., Phys. Rev. D **84**, 114010 (2011); Khodjamirian A et al., Eur. Phys. J A **48**, 31 (2012)
- [15] Grape S, *Studies on PWO Crystals and Simulations of the $\bar{p}p \rightarrow \bar{\Lambda}\Lambda, \bar{\Lambda}\Sigma^0$ Reactions for the PANDA experiment* (Ph. D. Thesis, Uppsala University, 2009)
- [16] Ikegami Andersson W K, *private communication*